

AD A 0 56788



LEVEL




CENTER FOR
PLASMA PHYSICS
AND
FUSION ENGINEERING

UNIVERSITY OF CALIFORNIA LOS ANGELES



his document has been approved or public release and sale; its istribution is unlimited.

029



EXCITATION OF LARGE AMPLITUDE PLASMA WAVES
BY RUNAWAY ELECTRONS

by

J. N. Leboeuf, T. Tajima, and G. J. Morales

PPG-349

April, 1978



Department of Physics
University of California
Los Angeles, California 90024

This document has been approved for public release and sale; is distribution is unlimited.

AD No. -ODC FILE COPY

78 07 24 029


J. N./Leboeuf, T./Tajima, G. J./Morales	KEPURIU	OCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FOR
EXCITATION OF LARGE AMPLITUDE PLASMA WAVES BY RUNAWAY ELECTRONS 7. AUTHOR(a) 1. N./Leboeuf, T./Tajima, G. J./Morales 8. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics University of California Los Angeles, California ONR Physics Program Office Arlington, Virginia 22217 13. MONITORING AGENCY NAME & ADDRESS(il diliterant from Controlling Office) Approved for public release; distribution unlimited. 16. DISTRIBUTION STATEMENT (a) the abstract entered in Block 20, Il diliterant from Report) Nonlinear waves, Runaway electrons 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side II necessary and Identity by Mack number) Nonlinear waves, Runaway electrons 18. ABSTRACT (Continue on reverse side II necessary and Identity by Mack number) A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electroic field enhances the excitation of plasma waves through runaway electroic field enhances the excitation of plasma waves through runaway electroic field enhances the excitation of plasma waves through runaway electroic field enhances a longated tail in the distribution function gives rise tacoustic-type modes which interact strongly with the runaway population. To waves produce a strong drag which reduces the runaway momentum increase from 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	1. REPORT NUMBER	2. GOVT ACCES	
EXCITATION OF LARGE AMPLITUDE PLASMA MAVES BY RUNAMAY ELECTRONS AUTHORY J. N./Leboeuf, T./Tajima, G. J./Morales PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics / University of California 90024 1. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 2. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office) Approved for public release; distribution unlimited. 5. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II different from Report) Nonlinear waves, Runaway electrons 6. SUPPLEMENTARY NOTES 6. SUPPLEMENTARY NOTES 6. ABSTRACT (Continue on reverse side II necessary and Identify by block number) A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electroric from Program and Program of Program Program and Program			19)
AUTHOR(a) J. N./Leboeuf, T./Tajima, G. J. Morales J. N./Leboeuf, T./Tajima, G. J. Morales PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics / University of California Los Angeles, California 90024 1. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 4. MONITORING AGENCY NAME & ADDRESS(il diliterent from Controlling Office) Approved for public release; distribution unlimited. 5. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, Il diliterent from Report) Nonlinear waves, Runaway electrons 6. ABSTRACT (Continue on reverse side if necessary and identify by block number) Nonlinear waves, Runaway electrons 6. ABSTRACT (Continue on reverse side if necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electroic field is electric field enhances the excitation of plasma waves through runaway electroic field is electric field enhances the excitation of plasma waves through runaway electroic field is electric field on the distribution function gives rise tacoustic-type modes which interact strongly with the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	TITLE (and Subtitle)		TYPE OF REPORT & PERIOD COV
J. N./Leboeuf, T./Tajima, G. J. Morales PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics J. University of California Los Angeles, California 90024 1. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 4. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office) 5. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 6. DISTRIBUTION STATEMENT (of this abstract entered in Block 20, II dillerent from Report) Nonlinear waves, Runaway electrons 6. ABSTRACT (Continue on reverse side If necessary and Identify by block number) Nonlinear waves, Runaway electrons 6. ABSTRACT (Continue on reverse side If necessary and Identify by block number) A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electror from presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. Twaves produce a strong drag which reduces the runaway momentum increase from a threshold DC electric field is found			BY Technical replay
J. N./Leboeuf, T./Tajima, G. J./Morales Department of Physics / Department of Physics / Department of Physics / Department of Physics / University of California goodal Los Angeles, California goodal Los Angeles, California goodal normal physics Program Office Arlington, Virginia 22217 Monitoring Agency Name a Address (It different from Controlling Office) Approved for public release; distribution unlimited. DISTRIBUTION STATEMENT (of the abstract entered in Block 30, If different from Report) Approved for public release; distribution unlimited. E. DISTRIBUTION STATEMENT (of the abstract entered in Block 30, If different from Report) Nonlinear waves, Runaway electrons A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electror from the presence of an elongated tail in the distribution function gives rise tacoustic-type modes which interact strongly with the runaway population. Twaves produce a strong drag which reduces the runaway momentum increase from 2/3 to 1/2 of the free fail value. A threshold DC electric field is found	RUNAWAY ELECTRONS	•	6. PERPORUTE DASCREPORT NUMBER
J. N./Leboeuf, T./Tajima, G. J. Morales Performing organization name and address Department of Physics J University of California Los Angeles, California 90024 1. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 13. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office) 14. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office) 15. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, If different from Report) Nonlinear waves, Runaway electrons 16. Supplementary notes 17. ABSTRACT (Continue on reverse elds if necessary and Identify by block number) Nonlinear waves, Runaway electrons 18. ABSTRACT (Continue on reverse elds if necessary and Identify by block number) A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrops repersured as a coustic-type modes which interact strongly with the runaway population. Twaves produce a strong drag which reduces the runaway momentum increase from 2/3 to 1/2 of the free fall value. A threshold DC electric field is found			
D. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics / University of California Los Angeles, California 90024 1. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 13. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office) Approved for public release; distribution unlimited. 15. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II dillerent from Report) Nonlinear waves, Runaway electrons 16. Supplementary notes 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II dillerent from Report) Nonlinear waves, Runaway electrons 18. ABSTRACT (Continue on reverse side II necessary and Identify by block number) A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electron from Program and Program and Identify by block number) A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electron gives rise the acoustic-type modes which interact strongly with the runaway population. The waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	- AUTHOR(s)	Company of the Compan	(I)
University of California 90024 1. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 13. NUMBER OF PART 23 14. MONITORING AGENCY NAME & ADDRESS(II diliterent from Controlling Office) Approved for public release; distribution unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II diliterent from Report) Nonlinear waves, Runaway electrons 8. KEY WORDS (Continue on reverse elde II necessary and Identify by block number) Nonlinear computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electroic field enhances the excitation of plasma waves through runaway electroic field enhances the excitation of plasma waves through runaway electroic field enhances the excitation of plasma waves through runaway electroic field enhances the excitation of plasma waves through runaway electroic field enhances the tacoustic-type modes which interact strongly with the runaway population. Twaves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	J. N./Leboeuf, T./	Tajima, • G. J. Morales	
University of California Los Angeles, California 1. Controlling Office NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 14. MONITORING AGENCY NAME a ADDRESS(II different from Controlling Office) Approved for public release; distribution unlimited. 15. DECLASSIFICATION/DOWNGRAE SCHEDULE The Declassification/downgrae Schedule The Declassification/downgrae Schedule 16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, If different from Report) Nonlinear waves, Runaway electrons 16. ABSTRACT (Continue on reverse side if necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a Declassific computer simulation demonstrates that the application of an electric field enhances the excitation of plasma waves through runaway electrons free presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. Twaves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found			10. PROGRAM ELEMENT, PROJECT, 1
Los Angeles, California 90024 11. CONTROLLING OFFICE NAME AND ADDRESS ONR Physics Program Office Arlington, Virginia 22217 12. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office) Approved for public release; distribution unlimited. 15. DESCLASSIFICATION/DOWNGRAE SCHEDULE 16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II different from Report) Nonlinear waves, Runaway electrons 16. Supplementary notes 17. Distribution on reverse elde II necessary and Identify by block number) Nonlinear waves, Runaway electrons 18. ABSTRACT (Continue on reverse elde II necessary and Identify by block number) A relativistic computer simulation demonstrates that the application of a Delectric field enhances the excitation of plasma waves through runaway electron from presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. Tweeton of the free fall value. A threshold DC electric field is found at the strong in the presence of the free fall value. A threshold DC electric field is found increase from 2/3 to 1/2 of the free fall value. A threshold DC electric field is found			
ONR Physics Program Office Arlington, Virginia 22217 14. MONITORING AGENCY NAME & ADDRESS(II diliterent from Controlling Office) 15. SECURITY CLASS. (SI this report) Approved for public release; distribution unlimited. 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II different from Report) Nonlinear waves, Runaway electrons 18. Supplementary notes 19. KEY WORDS (Continue on reverse side II necessary and identify by block number) Nonlinear waves, Runaway electrons 19. ABSTRACT (Continue on reverse side II necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrons The presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. To waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found			
Physics Program Office Arlington, Virginia 22217 18. MONITORING AGENCY NAME & ADDRESS(II diliterent from Controlling Office) 18. SECURITY CLASS (III for a poor) 18. DECLASSIFICATION/DOWNGRAE 18. DECLASSIFICATION/DOWNGRAE 18. DECLASSIFICATION/DOWNGRAE 18. DECLASSIFICATION/DOWNGRAE 18. DECLASSIFICATION/DOWNGRAE 18. DECLASSIFICATION/DOWNGRAE 19. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II diliterent from Report) 19. NORTHER WORDS (Continue on reverse side if necessary and identify by block number) Nonlinear waves, Runaway electrons 19. ABSTRACT (Continue on reverse side if necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electron presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found		ME AND ADDRESS	
Arlington, Virginia 22217 MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office) 18. SECURITY CLASS (II this report) Approved for public release; distribution unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, II dillerent from Report) Nonlinear waves, Runaway electrons A RESTRACT (Continue on reverse side if necessary and identify by block number) A relativistic computer's simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrons the presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found		ffice	
Approved for public release; distribution unlimited. 77. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different from Report) 18. SUPPLEMENTARY NOTES 18. SUPPLEMENTARY NOTES 18. ABSTRACT (Continue on reverse side if necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrons the presence of an elongated tail in the distribution function gives rise t acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found			HARDS (12)23
Approved for public release; distribution unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different from Report) 8. SUPPLEMENTARY NOTES 9. KEY WORDS (Continue on reverse side if necessary and identify by block number) Nonlinear waves, Runaway electrons A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrope presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. The waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	4. MONITORING AGENCY NAM	E & ADDRESS(if different from Controlling	Office) 18. SECURITY CLASS. (of this report)
Approved for public release; distribution unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different from Report) 8. SUPPLEMENTARY NOTES 9. KEY WORDS (Continue on reverse side if necessary and identify by block number) Nonlinear waves, Runaway electrons 1. ABSTRACT (Continue on reverse side if necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrope produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found			15a. DECLASSIFICATION/DOWNGRAD
Approved for public release; distribution unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 8. SUPPLEMENTARY NOTES 9. KEY WORDS (Continue on reverse elde if necessary and identify by block number) Nonlinear waves, Runaway electrons 1. ABSTRACT (Continue on reverse elde if necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrons the presence of an elongated tail in the distribution function gives rise that acoustic-type modes which interact strongly with the runaway population. To waves produce a strong drag which reduces the runaway momentum increase from 2/3 to 1/2 of the free fall value. A threshold DC electric field is found			SCHEDULE
Nonlinear waves, Runaway electrons ABSTRACT (Continue on reverse elde if necessary and identify by block number) A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway electrone presence of an elongated tail in the distribution function gives rise t acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	Approved for publ		unlimited.
Nonlinear waves, Runaway electrons A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway elec The presence of an elongated tail in the distribution function gives rise t acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found		ic release; distribution	
A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway elec The presence of an elongated tail in the distribution function gives rise t acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	7. DISTRIBUTION STATEMENT	ic release; distribution	
A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway elec The presence of an elongated tail in the distribution function gives rise t acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	7. DISTRIBUTION STATEMENT	ic release; distribution	llerent from Report)
A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway elec The presence of an elongated tail in the distribution function gives rise t acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	7. DISTRIBUTION STATEMENT 18. SUPPLEMENTARY NOTES 9. KEY WORDS (Continue on rev	ic release; distribution (of the abetract entered in Block 20, if di	llerent from Report)
A relativistic computer simulation demonstrates that the application of a D electric field enhances the excitation of plasma waves through runaway elec The presence of an elongated tail in the distribution function gives rise t acoustic-type modes which interact strongly with the runaway population. T waves produce a strong drag which reduces the runaway momentum increase fro 2/3 to 1/2 of the free fall value. A threshold DC electric field is found	17. DISTRIBUTION STATEMENT 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on rev	ic release; distribution (of the abetract entered in Block 20, if di	llerent from Report)
above which there is no collective slowdown.	17. DISTRIBUTION STATEMENT 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on rev	ic release; distribution (of the abetract entered in Block 20, if di	llerent from Report)
	19. KEY WORDS (Continue on rev Nonlinear wave: A relativistic comprelectric field enhal The presence of an acoustic-type modes waves produce a streacy 2/3 to 1/2 of the f	ic release; distribution (of the abetract entered in Block 20, if di error elde if necessary and identify by block s, Runaway electrons error elde if necessary and identify by block uter simulation demonstrat nces the excitation of pla elongated tail in the dist which interact strongly w ong drag which reduces the ree fall value. A thresho	trumber) the number of a Discount of a Discount function of a Discount

472 267 401 733

hc

ABSTRACT

A relativistic computer simulation demonstrates that the application of a DC electric field enhances the excitation of plasma waves through runaway electrons. The presence of an elongated tail in the distribution function gives rise to acoustic-type modes which interact strongly with the runaway population. The waves produce a strong drag which reduces the runaway momentum increase from 2/3 to 1/2 of the free fall value. A threshold DC electric field is found above which there is no collective slowdown.



The evolution of runaway electrons taking into consideration the selfconsistent excitation of large amplitude collective oscillations is of considerable interest1,2,3 at the present time. This interest has been stimulated by recent experimental observations4,5,6 in tokamak devices of runaway electrons accompanied by the spontaneous emission of intense radiation in various4,5 frequency intervals. The study of runaway electrons provides a suitable arena to test and develop various theories of collective interactions in plasmas whose applicability can extend beyond just the runaway problem. Recent theoretical studies rely on assumptions concerning a flat momentum distribution parallel to the applied electric field. However, many details associated with the wave-particle interactions that generate and alter such distributions still remain to be illustrated. One approach previously used to examine some of the relevant physics has been the Fokker-Planck type of simulation^{7,8} in which the wave-particle interactions are assumed to be described by quasilinear theory. The latter assumption is also characteristic of most of the theoretical treatments of the problem. In yet another work, non-relativistic particle simulations without DC field and starting from a flat distribution have been performed8 to verify that collective effects could lead to a positive slope.

This study examines some of the wave-particle interactions and the nature of the collective modes present in the evolution of runaway electrons, employing a full particle computer simulation. The major effects found are:

(1) the runaway electrons exhibit large excursions in momentum space not amenable to a quasilinear discription, (2) the momentum distribution function develops an elongated tail which gives rise to modes with frequencies lower than the plasma frequency, 9 (3) the excitation of large amplitude waves reduces the rate of increase of the average runaway momentum from 1/2 to 2/3 of the free fall value.

A one dimensional relativistic electrostatic particle code is used. The system is initialized with a low density cold electron beam and a Maxwellian background distribution. The external DC electric field \mathbf{E}_{DC} is applied selectively to fast particles with momentum above a set value, chosen smaller than the initial beam momentum and larger than the initial average momentum associated with the thermal velocity of the background particles v_{TO}. This technique models the effect of collisions which would prevent the slower particles from running away in a real system. In the runs investigated we have considered two different values of the beam to background density ratio, $n_b/n_0 = 1/9$ and 1/180. The system length is 256 λ_{DO} , where λ_{DO} is the Debye length at t = 0; 256 beam electrons and 2304 bulk electrons are followed. Appropriate measures are taken to handle the discreteness of the beam electrons. Typical parameter values are as follows. The beam momentum at t = 0 is $p_b = 7.63 \text{ mv}_{T0}$, where m is the electron mass. The limiting value of the momentum above which E_{DC} is applied is 6.0 mv_{T0}. The speed of light c is taken as 9.0 v_{T0} , and $E_{DC} = \frac{eE_{DC}}{m\omega_p v_{T0}} = 0.1$, where ω_p is the electron plasma frequency and e the electron charge. Each electron is represented by a Gaussian-shaped finite size particle of width 1.0 λ_{DO} . A typical run extends to t * $10^3 \omega_n^{-1}$.

The early evolution of the system is dominated by the two-stream instability triggered by the low density runaway beam as shown in Fig. 1(a). During the early phase one observes the exponential growth characteristic of such an instability. Unlike the E_{DC} = 0 case, the instability does not saturate by particle trapping 10 , but rather the wave energy grows secularly eventually reaching a value roughly 10 times larger than the non-runaway beam saturation level, as illustrated in Fig.1(a). This figure exhibits the time evolution of the total wave energy for n_b/n_o = 1/180; the continuous

curve corresponds to \tilde{E}_{DC} = 0.1 and the smaller dotted curve to E_{DC} = 0. The latter curve is the well-known trapped particle saturation pattern. The reason for the enhanced excitation of plasma waves is that the free energy of the beam continues to be replenished through the work done by the DC field, hence the beam nonlinearities act as a converter of DC to AC energy. Together with the secular increase in the total wave energy one observes that the wave spectrum cascades toward smaller wavenumbers k eventually reaching k < ω_{D}/c .

An examination of phase space (not shown) indicates that individual runaway electrons experience large momentum oscillations due to the interaction with the large amplitude waves. In this manner the spatially averaged momentum distribution f(p) develops an elongated tail which proceeds to expand in time toward large momenta. An example of such a distribution is shown in Fig. 1(b) for the case $n_b/n_0 = 1/9$ and at $\omega_p t = 550$. As seen in Fig. 1(b), the elongated tail is not smooth, but rather it exhibits the transient formation and subsequent destruction of small bumps (beams). These bumps, which act as converters of DC energy into wave energy, produce the rapid growth and damping of the wave energy seen in Fig. 1(a) over the interval $\omega_p t > 500$. The measured relaxation oscillation period is $56 \omega_p^{-1}$ for $n_b/n_0 = 1/180$. This relaxation period may be evaluated as follows. The Landau growth rate is given by

$$\gamma_L = \pi/2 \sum_{k} \frac{\omega_k^3}{k^2} \frac{\partial f(p)}{\partial p} \middle| v = \omega_k/k$$
 (1)

Here $\partial f/\partial p$ may be estimated as $n_R/(p_b p_M n_o)$, where p_M , the maximum momentum cut off, p_b , the effective gradient length of the bump, and n_R/n_o are 90 mv $_{TO}$, 3mv $_{TO}$ and 0.016 respectively from the simulations. Here n_R is the number of runaway electrons.

The summation is taken over bumps in resonance with the dominant components of the wave spectrum (modes 4 and 5), which yields $\gamma_L \sim 1 \times 10^{-2} \omega_p$. A much smaller value would be obtained from the overall (negative) slope of the runaway tail. Since the wave energy varies by 40% over the trough value, this leads to a relaxation oscillation period $\tau_L \sim 2 \frac{\ln 1.4}{\gamma_L} = 54 \omega_p^{-1}$. In the case $n_b/n_o = 1/9$, the measured relaxation oscillation period is $24 \omega_p^{-1}$, while an argument similar to the above yields $\tau_L \sim 21 \omega_p^{-1}$. The relaxation oscillation period then increases as the beam to plasma density ratio is lowered, as expected from the growth rate expression, Eq.(1). The above estimates also imply that an argument based on a steady state picture might be inadequate.

An assumption which permeates the quasilinear description of runaway behavior is that the real part of the dispersion relation is determined by the background (slow) particles. However, as seen in Fig. 1(b), the self-consistent distribution consists of an elongated tail. The dispersion relation for such a distribution may be written in the highly relativistic limit as

$$1 - \frac{\omega_{p}^{2}(1-\eta)}{\omega^{2}-3k^{2}v_{TO}^{2}} - \frac{mc}{p_{M}} \frac{\eta\omega_{p}^{2}}{\omega(\omega-kc)} = 0, \qquad (2)$$

where $\eta = n_R/(n_0 + n_R)$ and p_M is the maximum (cut-off) momentum, $p_M >> mc$. The modes thus introduced may be called an electroacoustic branch due to the hot electron population (elongated tail). The appearance of a dispersionless low frequency branch is indicative of the Van Kampen effect, since the cut-offs in runaway electron momenta (at 0 and p_M) introduce the noncancellation of the phases of the Van Kampen modes. 12 In Fig. 2 we compare the prediction of Eq.(2) with $p_M/mc = 13.3$ and $\eta = 0.016$ (solid curve) with the simulation results (solid dots) obtained by standard correlation methods. It is clearly 13 seen that the

elongated tail strongly modifies the wave dispersion, as expected. Furthermore, the strong nonlinearity produced by the large amplitude waves gives rise to multiple-harmonic branches (crosses)!⁴ A salient feature in Fig. 2 is that the system supports modes whose phase velocity is slower than the edge velocity of the expanding distribution for $k < \omega_p/c$. Consequently, the runaway particles are always in resonance with a collective mode. The excitation of such modes retards the free fall expansion of the runaway tail.

Although the interaction of an individual runaway electron with the large amplitude waves is quite complicated, the average behavior of the runaway population is remarkably simple, as shown in Fig. 3. The average runaway momentum \overline{p} (i.e., average above the cut-off at 6.0 mv_{T0}) is found to increase linearly with time, $\overline{p} = p_0 + \alpha e E_{DC} t$, where the coefficient $\alpha \approx 2/3$ for $n_b/n_0 = 1/9$ and $\alpha \approx 1/2$ for $n_b/n_0 = 1/180$. This behavior can be understood through a simple scaling argument by realizing that for $n_b/n_0 = 1/9$ the number of runaway electrons is increasing only slightly and that the time averaged distribution function approximately has a self-similar triangular form given by

$$f(p) = \frac{2}{3\overline{p}} (1 - \frac{p}{3\overline{p}})$$
 $(0 \le p \le 3\overline{p})$ (3)

Conservation of momentum, $(d/dt)(p^2/2) = eEp$, with

$$(d/dt)(p^2/2) \equiv (d/dt) [(1/2) \int_0^{3\overline{p}} f(p)p^2dp] = (3/2) \overline{p} \frac{a}{p}$$
 (4)

and

$$eEp \equiv e \int_{0}^{3\overline{p}} f(p) p \cdot E_{DC} dp = eE_{DC} \overline{p}$$
 (5)

yields

$$\frac{\cdot}{p} = (2/3) eE_{DC}$$
 (6)

in excellent agreement with the simulations (see Fig. 3).

In addition to the classical runaway threshold (i.e. the Dreicer field¹⁵), we find another threshold DC field beyond which the electrons keep running away toward infinite momenta without any strong collective phenomena taking place. This critical DC field, $E_{\rm cr}$, is determined from the peak wave amplitude¹⁶ excited by a beam when $E_{\rm DC}$ = 0, i.e.

$$E_{cr} = \left[4\pi n_b \gamma_0 mc^2 \ s(1+s)^{-5/2}\right]^{\frac{1}{2}} \tag{7}$$

where γ_0 is the relativistic factor and $s = \beta^2 \gamma_0 (n_b/2n_o)^{1/3}$, $\gamma_0 = (1 - \beta^2)^{-\frac{1}{2}}$. When $E_{DC} > E_{cr}$, the waves fail to sustain oscillations and thus slow down the particles. In the case $n_b/n_o = 1/9$, a strong two-stream instability is still experienced in the simulation for $\tilde{E}_{DC} = 1.0$, while no two-stream instability nor any strong wave-particle interactions occur for $\tilde{E}_{DC} = 2.0$. In fact, it is observed that as E_{DC} approaches E_{cr} the level of AC fluctuations dramatically increases and then for $E_{DC} > E_{cr}$ it remains at the thermal level. Simulation values of $\gamma_0 = 1.31$, $c = 9.0 \ v_{TO}$, $n_b/n_o = 1/9$ and s = 0.21 (at t = 0) in Eq.(7) yield $\tilde{E}_{cr} \simeq 1.25$, confirming the simulation results. For $n_b/n_o = 1/180$, this threshold is found to lie between $\tilde{E}_{DC} = 0.1$ and $\tilde{E}_{DC} = 0.5$ in the simulation, while Eq.(7) predicts $\tilde{E}_{cr} = 0.3$.

The physical picture which emerges out of this investigation is the following. The presence of a DC electric field $E_{\rm DC}$ < $E_{\rm cr}$ prevents the usual saturation of the beam-plasma instability. Consequently, large amplitude plasma waves are excited and cause large momentum fluctuations in the runaway population. The strong nonlocal mixing in momentum space produces an elongated momentum distribution which supports Van Kampen (electroacoustic) type of modes. The self-consistent excitation of these modes gives rise to the transient growth and decay of bumps in the distribution function accompanied by relaxation oscillations in the wave energy. The drag produced by these waves reduces the runaway momentum increase below the free fall value.

ACKNOWLEDGMENT

The authors acknowledge fruitful discussions with Professor

J. M. Dawson. This research was supported by the National Science

Foundation Contract PHY 76-83686, the Department of Energy Contract

EY-76-C-03-0010 PA26/TASK III, and the Office of Naval Research Contract

N00014-75-C-0476 P00004.


REFERENCES

- 1. K. Molvig, M. S. Tekula, and A. Bers, Phys. Rev. Lett. 38, 1404 (1977).
- 2. C. S. Liu and Y. Mok, Phys. Rev. Lett. 38, 162 (1977).
- 3. C. S. Liu, Y. C. Mok, K. Papadopoulos, F. Engelman, and M. Bornatici, Phys. Rev. Lett. 39, 701 (1977).
- 4. A. A. M. Oomens, L. Th. M. Ornstein, R. R. Parker, F. C. Schuller, and R. J. Taylor, Phys. Rev. Lett. 36, 255 (1976)
- D. A. Boyd, F. J. Stauffer, and A. W. Trivelpiece, Phys. Rev. Lett. <u>37</u>, 98 (1976).
- 6. A. E. Costley and TFR Group, Phys. Rev. Lett. 38, 1471 (1977).
- 7. R. M. Kulsrud, Y.-C. Sun, N. K. Winsor, and H. A. Fallon, Phys. Rev. Lett. 31, 690 (1973).
- B. Basu, B. Coppi, K. Molvig, F. Pegoraro, I. Haber, B. Hui, P. Palmadesso,
 K. Papadopoulos, and N. Winsor in "Proceedings of the International
 Conference on Plasma Physics and Controlled Nuclear Fusion Research", 1976
 (International Atomic Energy Agency, Vienna, Austria, 1977), Vol. II, p.455.
- 9. R. Shanny, W. L. Kruer, and J. M. Dawson, in "Proceedings of the Second Conference on Numerical Simulation of Plasma", Los Alamos, New Mexico, 1968, Paper A2 (unpublished).
- 10. S. Kainer, J. M. Dawson, and T. Coffey, Phys. Fluids 15, 2419 (1972).
- 11. A multibeam distribution function has recently been observed experimentally in a toroidal device. K. W. Gentle, G. Leifeste, and R. Richardson, Phys. Rev. Lett. 40, 317 (1978).
- 12. N. G. Van Kampen, Physica 21, 949 (1955).
- 13. Because the fluctuating cut-off velocities mix up the phases of the different wavenumbers, the electroacoustic branch has not been clearly observed in nonrelativistic simulations (e.g. Ref. 9). Here, however, cut-off velocities are already close to c.

- 14. The relative intensities of the harmonics are a steeply decreasing function of wavenumber. Unless the waves are strongly excited, as is the case here, a clear mapping of the multiple harmonic branches might be difficult.
- 15. H. Dreicer, Phys. Rev. 115, 238 (1959).
- 16. L. E. Thode and R. N. Sudan, Phys. Rev. Lett. 30, 732 (1973).

FIGURE CAPTIONS

- FIG. 1 (a) Time evolution of the wave energy in the presence of runaway electrons. With $\tilde{E}_{DC} = 0.1$ (solid curve), the AC wave energy is sustained to a much higher level than with $E_{DC} = 0$ (dotted curve). $n_b/n_o = 1/180$. (b) The time averaged momentum distribution function at $t = 550 \omega_p^{-1}$ shows the elongated tail. $n_b/n_o = 1/9$.
- FIG. 2 Electroacoustic dispersion relation measured (solid dots) and predicted by theory (solid curve) for $n_b/n_o = 1/180$. Note that there appear second, third and fourth harmonic branches (crosses) in the dispersion relation. The spectral intensity of the harmonic modes is highest at the crossing of the harmonic and electroacoustic branches.
- FIG. 3 Evolution of the average runaway momentum. The solid dots represent the simulation results, the dashed line $\overline{p} = p_0 + 2/3 \text{ eE}_{DC} t.$ The average runaway momentum first follows the free fall line and eventually asymptotes to 2/3 of the free fall value.






FIGURE 2




FIGURE 3

JELA PLASMA PHYSICS CROUP REPORTS

- "Propagation of ion Acoustic Waves Along Cylindrical Plasma Columns", Phys. Fluids 9, 1261 (1966)(A.Y. Wong). "Stability Limits for Longitudinal Waves in Ion Beam-Plasma Interaction", Phys. Fluids 9, 1064 (1966)(B.D. Fried and
 - Tried and S.L. Ossabon).
 Free and S.L. Ossabon).
 Special and S.L. Ossabon).
 Special and S.L. Ossabon of a Calissional Electron Plassa", Phys. Fluids 9, 292 (1966)(8.D. Fried, A.N. Kaufman L. Secks).
- PPG-75 PPG-76 PPG-77 "inferent of Colisions on Electrostatic los Cyclotron Mayor", Phys. Lett., 21, 157 (1966)(A.Y. Wong, D. Judd and F. Hal).
 "Interestical Research in Bassa and Plasmas." Ball. As. Phys. Soc., 1152 (1965)(A.Y. Wong, D. Judd and F. Mail.).
 "Interestical Research Echoes from a Highly Ionized Plasma", Phys. Rev. Lett., 21, 157 (1964)(D.E. Kaplan and R.H. Hill
 "Excitation and Damping of Differ Mayors, Rev. Lett., 18, 256 (1965)(A. Mong, P. Rev. Colory).
 "The Guidding Center Approximation is Least Order", J. Flesse Phys., 1, 305 (1965)(A. Mong, J. J.
 "Plasma Streaming into a Magnetic Field", Dissertation (S.L. Orsakon) November (1965).
 "Noperative Effects in Plasma Echo Phenomena", Proc. of 1st Int'l. Conf. on Quicecont Ionsan, Francisi, Rome, 1967,
 "Annual Mechanical Study of the Electron Get Vist for Particle Method" Dissertation, March (1967)(M.E. Memain).
 "Linear and Menlinear Theory of Grid Encitation of Low Frequency Neves in a Plasmas", Dissertation, April (1967) (G.L. P.
 - PPG-81
- PPG-83 usion and Diffusion of an Isolated Plasma Column", Dissertation, May(1967)(J. Myman). Approximation for the Fissem Dispersion Function", Phys. Fluids <u>11</u>, 249 (1968)(B.D. Fried, C.L. Hedrick and

- Proc. 18 "Experimental lowestigation of Electron Research Peaceness", Dissertation, August (1967)(J.S. DeGroot).

 Proc. 18 "Experimental lowestigation of Electron Research Proc. 20 Zed Inc. 1. Symposium on Fluctuations and Diffusion in Plasmas, June Proc. 20 Test Rev. Compiled Revenes Proc. 20 Zed Lett. 2, 20 Zed Let
- PPG-104 PPG-105 PPC-46 "A PPG-47
 - In system of Conflictor The Conductivity Tensor of a Collisionless Plasma in a Magnetic Field", Fastschrift for Gregor Waters (10 of Chicago Perms, 1809) March (1809) (L.). Fried and C. Medrich).

 Est Mensurements of Linear Growth Asses and Monitare Statestic Coefficients", Phys. Rev. Lett., 23, 165 (1969)

 (A.Y. Mange and F. Mal.).

 Series Precipitation Polisations", J. Geophys. Res., 75, 1779 (1970) (F. Coroniti and C.F. Kennal).

 West of Febbra-Flanck Collisions on Plasma March Endows, Phys. Relief 19, 186 (1970) (G. Johnston).

 Series Abstract Theory of Graft Entitation of Low Frequency Naves in a Plasma", July (1969) (G. Johnston).

 HAY Goldsman).

 (B.Y. Goldsman).
- for the first of t
 - or Strong inchessatic News New Interaction", January (1979)(R.J. Taylor and H. Heatl).

 In a Distribution is large two and Econow', Phys. Finds 19, 3749 (1970)(R. Heat and R.J. Taylor).

 on of los Cycleton Numeric News, Phys. Finds 19, 2774 (1970)(R. Anit and H. Heatl).

 on of los Cycleton Numeric News, Phys. Finds 19, 2774 (1970)(R. Anit and H. Jean).

 In a Mangachet Properties of the Plans Dispersion Nuclearing, Physical 1970)(R. Mone and G. Johnston).

 In a Los California and Los Numer Translates States", June 1979) (G. Moter).

 In a Descripting of Stanger (2011stonies States", J. Plans Phys. 4, 285 (1970)(F. V. Orrontti).

19-944

- "urbalmens Estructure of Einite Bergendicular Pers Social", J. Googhy, Rev., 25, 700 (1990)(R. Corenit). "Steppening of Lee Accessic Muses and Forestican of Collisionless Electrostatic Social", April (1970)(R. Teylest). "A Manhool of Studying Trapped Particles Mahavior in Magnetic Geometries", Phys. Rev. Lett., 25, 7702 (1970)(G.S. Liu
- PPG-74

PPG-86

- PRICATA AND STATEMENT OF STATEMENT OF THE WARREST CHARACTERS. THE AND STATEMENT OF PE-38 PE-93 PE-93 PE-93

PPC-94

PPG- 100

PPG-98 PPG-99 PPG-103

PPG-101

- (c.) Wong and D. L. Jassby).

 (d.) Wong and D. L. Jassby).

 (e.) Wong and D. L. Jassby).

 (f.) Wong and D. L. Jassby).

 (e.) Wong and D. L. Jassby).

 (f.) Fareball institute, Praccett, Tally, Spreamber, 1971 (Pleaus Press, Karl Schindler, 1972)(F.) Coronitis and C. F. Leans).

 (f.) Fareball instruction with Luars Appoint Faldy. J. Pleans Phys., November (1971)(C. Sisce and C. E. Leans).

 (f.) Fried and F. V. Cervalit).

 (h) Fried and F. V. Cervalit).

 (h) Fried and F. V. Cervalit.

 (h) Fried and Cervality and Certailed Nuclear Plane Fried F. V. Sephys. Res., 72, 214 (1977) (R. Seeriel and A. Wang M. Wenner, F. F. Kenner).

 (h) Fried and Schwarzeric Cervalities F. V. Kenner, 72, 4197 (1972) (K. Les. P. K. Kenner).

 (h) Fried and Schwarzeric Cervalities F. V. Cervalit.

 (h) Fried B. V. Cervalit.

 (h) Fried PPG-107 PPG-108 PPG-109
 - - PPC-111

 C. F. Korman J.

 C. F. Lyon and P. V. Corvaitt J.

 C. F. Lyon and P. V. Corvaitt J.

 C. F. Lyon and P. C. F. Korman J.

 C. F. Lyon and R. M. F. Korman J.

 C. F. Lyon and R. M. Thoman J.

 C. K. Lyon and R. M. Morne J.

 C. M. Washing and R. M. Thoman J.

 C. M. Washing and R. M. Thoman J.

 C. M. Washing A. K. W. Cornait and R. M. Thoman J.

 C. M. Washing A. K. W. Cornait and R. M. Thoman J.

 C. M. Washing A. K. W. Cornait and R. M. Thoman J.

 C. M. Washing A. K. W. Cornait and R. M. Thoman J.

 C. M. Washing A. K. W. Cornait and R. M. Thoman J.

 C. M. Washing A. K. W. Cornait and R. M. Thoman J.

 C. M. Washing A. K. W. Cornait and R. M. Thoman J.

 C. M. Washing A. M. W. Cornait and R. M. Thoman J.

 C. M. Washing A. M. W. Cornait and R. M. Thoman J.

 C. M. Washing A. M. W. Cornait and R. M. Thoman J

The state of the control of the cont

The state of the first interest and the control of the control of

J.M. Danton of Mightlys loss and Cneration of Intenso Neutral Boson", Applied Phys. Lett., 25, 579 (1974)(A.Y. Vong. J.M. Danton and W. Galellung. PR.- 183 "Prode

THIS PAGE IS BEST QUALITY PRACTICABLE FROM COPY FURNISHED TO DDC

A.T. Lin & J.M. Dawson, J. Comp. Phys. 23, 300 (1977).
"Relativistic Effects in Resonance Absorption", J.F. Drake and Y.C. Lee, Phys. of Fluids 19, 1772 (1976).
"Convective Electron Loss Cone Instabilities", M. Ashour-Abdalla and C.F. Kennel. Submitted to J. Geophys. Res. (1976). 81, 4531 (1976).

"A Self-Consistent Magnetostatic Particle Code for Numerical Simulation of Plasmas", J. Busnardo-Neto, P.L. Pritchett, "Energetic Radiation Belt Electron Precipitation; A Natural Depletion Mechanism for Stratospheric Ozone, R.M. Thorne, of Fluids, 19, 1975 (1976). "The Scattering of Cosmic Rays by Magnetic Bubbles", R.F. Flewelling & F.V. Coroniti, Astrophys. J., 205, 135 (1976). "San Francisco Abstracts - Papers to be Presented at San Francisco Meeting of the American Physical Society Division "The Earth's Magnetosphere", F.V. Coroniti, Accepted by Ency. of Phys. (1976). "Lossy Radial Diffusion of Relativistic Jovian Electrons", D.D. Barbosa and F.V. Coroniti, J. Geophys. Res. 81, 4553 "Nonlinear Interactions of Focused Resonance Cone Fields with Plasmas", R. Stenzel and W. Gekelman, Phys. Fluids 20, "Experiments on Parametric Instabilities in Laser-Beam Plasma Interactions", R. Chen, Presented at Nobel Symposium. "The Local Time Variation of ELF Emissions During Periods of Substorm Activity", R.M. Thorne, F.R. Church, "Parametric Instabilities with Finite Wavelength Pump", B.D. Fried, T. Ikemura, K. Nishikawa, and G. Schmidt, Phys. "The Effect of Pump Cutoff on Parametric Instabilities in an Inhomogeneous Plasma", J. VanDam and Y.C. Lee, (1976). "Relativistic Electrons and Whistlers in Jupiter's Magnetosphere", D.D. Barbosa and F.V. Coroniti, J. Geophys. Res. "Magnetic Field Reconnection in a Collisionless Plasma", F.V. Coroniti and A. Eviatar, Astrophys. J. Supp. 33, 189 W.J. Malloy, B.T. Tsurutani, J. of Geophys. Res., 82, 1585 (1977).
"An Investigation of Relativistic Electron Precipitation Events and their Association with Magnetospheric Substorm "Plasma Simulation on the CHI Microprocessor System", T. Kamimura, J.M. Dawson, B. Rosen, G.J. Culler, R.D. Levee Phys. of Fluids, 19, 1909 (1976). "Soliton-Like Structures in Plasmas", Y.C. Lee and G.J. Morales, March (1976). To appear in Rocky Mtn. Math. Jr. "Synchro-Compton Radiation Damping of Relativistically Strong Plasma Waves", C.F. Kennel, E. Asseo and R. Pellat, Activity", R.M. Thorne and T.R. Larsen, J. of Geophys. Res. 81, 5501 (1976).
"The Free Electron Laser", T. Kwan, J.M. Dawson & A.T. Lin, Phys. Fluids 20, 581 (1977).
"Stimulated Compton Scattering of Electromagnetic Waves off Plasma Ions", A.T. Lin and J.M. Dawson, Phys. Fluids Y.C. Lee, & G.J. Morales, Phys. Fluids 19, 849 (1976). "The Spiky Turbulence Generated by a Propagating Electrostatic Wave of Finite Spatial Extent", G.J. Morales, "Particle and Wave Dynamics in a Magnetized Plasma Subject to High RF Pressure", W. Gekelman & R.L. Stenzel, R. Thorne, Pure and Applied Geophys. 114, 497 (1976). "Free Boundaries for Plasmas in Surface Magnetic Field Configurations", B.D. Fried, J.W. VanDam & Y.C. Lee, "Parametric Excitation of Ion Density Fluctuations in the Relativistic Beam-Plasma Interaction", H. Schamel "Departures from Theory of the Experimental Line Profile of Helium I 4471.5A", J. Turechek, January (1976). "Maintenance of the Middle Latitude Nocturnal D-Layer by Energetic Electron Precipitation", W. Spjeldvik & "Stability of Shell Distributions", D.D. Barbosa, Submitted to Planetary and Space Sciences, July (1976). Science, 195, 287 (1977). "CTR Using the P-11B Reaction", J.M. Dawson, August (1976). Submitted to Astronomy and Astrophysics, July (1976). of Plasma Physics, November 15-19 (1976). 4 Y.C. Lee, Phys. Fluids 19, 690 (1976). Phys. Fluids 20, 1316 (1977). PPG-246 PPG-251 PPG-252 PPG-255 PPG-261 PPG-263 PPG-265 PPG-266 PPG-268 PPG-270 PFG-274 PPG-249 PPG-250 PPG-253 PPG-256 PPG-258 PPG-259 PPG-260 PPG-264 PPG-267 PPG-271 PPG-272 PPG-273 PPG-254 PPG-257 PG-247 PPG-248

THIS PAGE IS BEST QUALITY FRACTICABLE FROM COPY FURNISHED TO BUILD

B.D. Fried, Phys. Fluids, 20, 815 (1977). "Electromagnetic Interactions with Inhomogeneous Plasmas", A.Y. Wong. Submitted to Nuclear Fusion and RPI Oct. (1976). "The Importance of Electrostatic Ion-Cyclotron Instability for Quiet-Time Proton Auroral Precipitation", "Plasma Heating by a Relativistic Electron Beam with Secondary Instabilities", T. Tajima. Submitted to J. Plasma Phys. "Kinetic Theory of Tearing Instabilities", J.F. Drake & Y.C. Lee, Phys. Fluids 20, 1341 (1977).
"Jupiter's Magnetosphere", C.F. Kennel & F.V. Coroniti, Annual Rev. of Astro. & Astrophys., Submitted Nov., 1976.
"Direct Density Display with a Resonance Cone RF Probe", J. Ickovic, R.L. Stenzel & W. Gekelman, Rev. of Scientific "The Alfven-Ion Cyclotron Instability: Its Physical Mechanism and Observation in Computer Simulation", T. Tajima, K. Mima, and J.M. Dawson, Phys. Rev. Lett. 39, 201 (1977). "Phase Mixing in the Continuous Spectrum of Alfven Waves", P.L. Pritchett & J.M. Dawson, Phys. Fluids, Submitted "Particle Orbits and Loss Regions in UCLA Tokamaks", C.P. Lee, January (1977)
"Generation of Density Cavities and Localized Electric Fields in a Nonuniform Plasma", G.J. Morales & Y.C. Lee, Submitted to Phys. Fluids (1977). "Surface Multipole Guide Field for Plasma Injection", R.A. Breun, B.H. Rael, & A.Y. Wong, Review of Scientific "Higher Order Adiabatic Invariants and Non-Adiabatic Diffusion for Geometric Mirrors", Y.C. Lee, T.K. Samec, & "Low-Z Impurity Control for Reactor Size Tokamaks", R.J. Taylor and L. Oren.
"Computer Simulation of Current Penetration in a Plasma", A.T. Lin and J.M. Dawson, Submitted to Phys. Fluids, "Nonlinear Evolution of Collisionless and Semi-Collisional Tearing Modes", J.F. Drake and Y.C. Lee, Phys. Rev. M. Ashour-Abdalla & R.M. Thorne, Geophys. Res. Lett., 4, 45, (1977).
"Self Channeling of Whistler Waves", E.S. Weibel, Phys. Letts. 61, 37 (1977).
"Formation of Potential Double Layers in Plasmas", B.H. Quon & A.Y. Wong, Phys. Rev. Letts., 37, 1393 (1976).
"Plasma Confinement by Surface Magnetic Fields", T.K. Samec. "Possible Origins of Time Variability in Jupiter's Outer Magnetosphere", F.V. Coroniti & C.F. Kennel, Geophy. "Active Stimulation of Ionospheric Plasma", A.Y. Wong, April, 1977. "Confinement of High B Toroidal Plasmas and Currents in Surface Magnetic Fields", D. Mamas, R.W. Schumacher, "Langmuir Waves in a Spatially Periodic Potential", A. Baños, E.S. Weibel, and J.W. VanDam, Phys. Fluids, "Nonlinear Evolution of the Collisionless Tearing Mode", F.V. Coroniti, Phys. Rev. Lett. 38, 1355 (1977). Res. Lett., 4, 287 (1977). "Coupling of Lower-Hybrid Radiation at the Plasma Edge", G.J. Morales, Phys. Fluids, 20, 1164 (1977). "Model Analysis for Shear Stabilization of Collisionless Drift Waves", Y.C. Lee & T. Tange. "Surface Trappings of Oxygen in Tokamaks and its Removal", Lena Oren and R.J. Taylor. "The Ponderomotive Effect in a Magnetized Plasma", E.S. Weibel.
"The Fluid Theory of the Ponderomotive Effect in a Magnetized Plasma", E.S. Weibel. "Simulation Study of Bernstein Modes", T. Kamimura, T. Wagner & J.M. Dawson. 'Quasi-Linear Theory for Non-Uniform Turbulence", E.S. Weibel, June, 1977. "Super Conducting Surmacs", A. Wong and L. Miller, May (1977). A.Y. Wong and R.A. Breun, submitted to Phys. Rev. Lett. Phys. Fluids, 20, 1135 (1977). Instruments 48 (1977). instr. 48, 485 (1977). Lett. 39, 453 (1977). March, 1977. PPG-282 PPG-289 PPG-298 PPG-300 PPG-275 PG-276 PPG-278 PPG-284 PPG-285 PPG-286 PPG-287 PPG-292 PPG-294 PPG-296 PPG-297 PPG-301 PPG-302 PPG-303 PPG-281 PPG-291 PPG-293 PPG-304 PPG-277

"Ion Sound Turbulence in a Magnetoplasma", W. Gekelman and R.L. Stenzel, submitted to Phys. Fluids, March (1978). "Dynamics of Test Particles in a Turbulent Magnetoplasma", R.L. Stenzel and W. Gekelman, submitted to Phys. Fluids, submitted to Phys. Rev. Lett. April (1978). "Possible Origins of Time Variability in Jupiter's Outer Magnetosphere. 3. Variations in the Heavy Ion Plasma", submitted to J. of Comp. Phys., March, (1978). "Global Simulation of the Time-Dependent Magnetosphere", J.N. Leboeuf, T. Tajima, C.F. Kennel, and J.M. Dawson, "Excitation of Large Amplitude Plasma Waves by Runaway Electrons", J. N. Leboeuf, T. Tajima, and G. J. Morales, "Magnetic Field Generation by the Rayleigh-Taylor Instability," T. Tajima, K. Mima, and J.N. Leboeuf.

To be published in the Proceedings of Topical Meeting of Inertial Confinement Fusion (CLEOS). February, 1978. "Kinetic Theory of Ballooning Instabilities," Y.C. Lee and J.W. VanDam. To be published in the Proceedings of the High-Beta Tokamak Theory Workshop, Varenna, 1977 (by the U.S. DOE). February, 1978. "Excitation of Lower Hybrid Waves in a Finite Plasma," V.K. Decyk, J.M. Dawson, and G.J. Morales, submitted "Small Amplitude Waves in a Hot Relativistic Two Fluid Plasma," S. Hyun and C.F. Kennel, March (1978). "Nonlinear Evolution of Tearing Instabilities: Violations of Constant w," J.F. Drake, P.L. Pritchett, and "Splines and High Order Interpolations in Plasma Simulations", H. Okuda, A.T. Lin, C.C. Lin and J.M. Dawson "Parametric Instabilities and Ion Heating in a Two Ion-Species Plasma", C.C. Lin, A.T. Lin, and H. Okuda. "Penetration of a Plasma into a Linear Cusp", A. J. Lamm and K. F. MacKenzie, submitted to Phys. Fluids, "Kinetic Theory of m = 1 Internal Instabilities," J.F. Drake, submitted to Phys. Fluids, January, 1978. to Phys. Fluids. February, 1978. "Explosive Tearing Mode Reconnection in the Magnetospheric Tail," A.A. Galeev, F.V. Coroniti, and M. A. Eviatar, C. F. Kennel, and M. Neugebauer, accepted by Geophys. Res. Lett., April (1978). Y.C. Lee. Submitted to Phys. Rev. Letts., March (1978). "Safety of Fusion Reactors", Kastenberg, Okrent and Pomraning, October (1977). submitted to Geophys. Res. Lett, April (1978). Ashour-Abdalla, March (1978). March (1978). PPG-346 PPG-347 PPG-348 PPG-349 PPG-336 PPG-338 PPG-339 PPG-343 PPG-345 PPG-337 PPG-342 PPG-344 PPG-341